

Doxycycline for the Prevention of Bacterial Sexually Transmitted Infections

Julie Dombrowski, MD, MPH
Associate Professor, Department of Medicine, UW
Deputy Dir., HIV/STD Program, Public Health – Seattle & King County

July 25, 2019



Disclosures

Hologic (research support)

Outline

- The need
- The pros
- The cons
- My opinion and approach

The STATE of STDs in the United States



in 2017

THE NATION EXPERIENCES STEEP AND SUSTAINED STD INCREASES.



1.7 million

CASES OF CHLAMYDIA

22% increase since 2013



555,608

CASES OF GONORRHEA

67% increase since 2013



30,644

CASES OF SYPHILIS

76% increase since 2013

What Tools do we Have for STI Prevention?



Counseling & Education



Condom promotion



Frequent testing & treatment


Combination prevention



Safer Sex Plans HPV PrEP About En Español

Doxycycline Prophylaxis for STIs

Doxycycline post-exposure prophylaxis (PEP)




Doxycycline pre-exposure prophylaxis (PrEP)

```
doxy doxy time
```


Doxycycline PEP

- RCT in open label extension of IPERGAY (N=232)
 - Event-driven (2-1-1) HIV PrEP study in France
 - Doxy PEP vs. usual care
- Doxy 200mg x1 ~24h after sex (≤72h)
- Targeting CT & syphilis
- Median 7 pills/month



47% ↓in STI (69.7 vs 37.7/100 PY)

Doxycycline PrEP



Experience with Doxycycline Prophylaxis

- Widely used for a number of indications, including
 - Prevention of malaria, leptospirosis, Lyme
 - Treatment of moderate to severe acne
- Predictable and manageable side effects
 - Sun sensitivity
 - Esophagitis







Interest in Doxycycline PEP for STI among MSM

King County STD Clinic PrEP Program (N=35)


Atlanta, Birmingham, Chicago, NYC, SF, Seattle






African American & Latino MSM were more likely to report interest than non-Hispanic white MSM

Community Use?





- London sexual health clinic
- July 2018 survey
- 107 men on PrEP
- 8 (8%) had taken antibiotics to prevent STIs



Antimicrobial Resistance

- Antibiotic use = antibiotic resistance
- STI pathogens
 - N. gonorrhoeae doxycycline not used
 - T. pallidum no tetracycline resistance
 - C. trachomatis no tetracycline resistance
- Non-STI commensals/pathogens
 - S. aureus and S. pneumoniae
 - Commensal Neisseria
 - Enteric bacteria
- Broader microbiome effects?
- The unknown unknowns

Doxycycline Safety

Long-standing clinical experience = doxy is safe

but

- Recommended duration for acne: 12-16 weeks, beyond 24 weeks not recommended
- Weight gain (?)
- Benign intracranial hypertension rare association

Balance of Risks and Benefits?

Need to consider STD morbidity, not just infection



Where Are We Now?

- Need more evidence for effectiveness
 - ≥4 studies planned or ongoing internationally
- Insufficient evidence to recommend routine use in practice
- Pros
 - Doxycycline is generally safe
 - Patients are interested
 - Public health STD control: little else to offer
- Cons
 - Antimicrobial resistance
 - Uncertain balance of risks and benefits
- My approach: very rare and time-limited

Thank you



jdombrow@uw.edu



@DrJDombrowski